

Prevalence of *Chlamydia abortus* in Belgian ruminants

Prevalentie van Chlamydia abortus bij herkauwers in België

^{1,2}L. Yin, ¹K. Schautteet, ¹I.D. Kalmar, ³G. Bertels, ³E. Van Driessche, ⁴G. Czaplicki, ⁵N. Borel, ⁶D. Longbottom, ⁷D. Frélin, ⁷M. Dispas, ¹D. Vanrompay

¹Department of Molecular Biotechnology, Faculty of Bioscience Engineering, Ghent University, Coupure Links 653, B-9000 Ghent, Belgium

²College of Veterinary Medicine, Sichuan Agricultural University, Ya'an, 625014, China

³Dierengezondheidszorg Vlaanderen, Hagenbroeksesteenweg 167, B-2500 Lier, Belgium

⁴Association Régionale de Santé et l'Identification Animal, Avenue Alfred Deponthière 40, B-4431 Loncin, Belgium

⁵Institute of Veterinary Pathology, University of Zürich, Vetsuisse Faculty Zürich, Winterthurerstrasse 268, CH-8057 Zürich, Switzerland

⁶Moredun Research Institute, Pentlands Science Park, Bush Loan, Edinburgh EH26 0PZ, UK

⁷Department of General Bacteriology, Veterinary and Agrochemical Research Centre, Groeselenberg 99, B-1180 Ukkel, Belgium

Daisy.Vanrompay@ugent.be

ABSTRACT

Chlamydia (C.) abortus enzootic abortion still remains the most common cause of reproductive failure in sheep-breeding countries all over the world. *Chlamydia abortus* in cattle is predominantly associated with genital tract disease and mastitis. In this study, Belgian sheep (n=958), goats (n=48) and cattle (n=1849) were examined, using the ID Screen™ *Chlamydia abortus* indirect multi-species antibody ELISA. In the sheep, the highest prevalence rate was found in Limburg (4.05%). The animals of Antwerp, Brabant and Liège tested negative. The prevalence in the remaining five regions was low (0.24% to 2.74%). Of the nine goat herds, only one herd in Luxembourg was seropositive. In cattle, the highest prevalence rate was found in Walloon Brabant (4.23%). The animals of Limburg and Namur tested negative. The prevalence rate in the remaining seven regions ranged between 0.39% and 4.02%.

SAMENVATTING

Enzoötische abortus veroorzaakt door *Chlamydia (C.) abortus* is nog steeds de meest voorkomende oorzaak van reproductiestoornissen bij schapen. Bij runderen wordt *C. abortus* geassocieerd met infecties van het geslachtsstelsel en mastitis. In de voorliggende studie werden Belgische schapen (n=958), geiten (n=48) en runderen (n=1849) onderzocht aan de hand van de ID Screen™ *Chlamydia abortus* indirecte multispecies antistof ELISA. Bij de schapen werd de hoogste prevalentie gevonden in Limburg (4,05%). De dieren van Antwerpen, Brabant en Luik waren negatief. De prevalentie in de overige vijf regio's was laag (0,24% tot 2,74%). Van negen geitenboerderijen was slechts één bedrijf uit Luxemburg positief. Bij de runderen werd de hoogste seroprevalentie in Waals-Brabant (4,23%) gevonden. De dieren uit Limburg en Namen waren negatief. De prevalentie in de overige zeven regio's varieerde van 0,39% tot 4,02%.

INTRODUCTION

Chlamydiaceae are gram-negative obligate intracellular bacteria. Recently, the assignment from the single genus *Chlamydia* into two genera, *Chlamydia* and *Chlamydophila* by Everett et al. (1999) has been reconsidered. Based on comparative genome analysis of several *Chlamydiaceae* genomes, the *Chlamydia-*

ceae are currently again reunited into a single genus, *Chlamydia* (Kuo and Stephens, 2011).

Ruminants can become infected with *Chlamydia (C.) abortus*, *C. pecorum*, *C. psittaci* and rarely with *C. suis* (Reinhold et al., 2011). *Chlamydiaceae* infections in cattle (*Bos taurus*) have been known to occur since 1940, when McNutt isolated intracellular organisms from cases of sporadic bovine encephalomyelitis

in feedlot cattle (McNutt and Waller, 1940). Thereafter, a number of studies worldwide reported epizootic bovine abortion in cattle caused by *C. abortus*. The pathogen also caused bovine mastitis, epididymitis and seminal vesiculitis, and was excreted in bull semen (Kaltenboeck et al., 2005; Reinhold et al., 2011). Chlamydial strains from ruminant abortion were first classified as mammalian *Chlamydia psittaci* serotype 1, mammalian *C. psittaci* biovar 1 or *C. psittaci* outer membrane protein A (*ompA*) gene type B577 strains (Kaltenboeck et al., 1993). Later on, the organism was reclassified as *C. abortus* (Everett et al., 1999).

In the previous century, a second chlamydial agent was reported to be associated with growth retardation, abortion, sporadic bovine encephalomyelitis, pneumonia, enteritis, polyarthritis, keratoconjunctivitis, nephritis or purulent endometritis in cattle. This chlamydial agent was first identified as mammalian *C. psittaci* serotype 2 or mammalian *C. psittaci* biovar 2. However, it became clear that these disease manifestations were actually induced by a serologically and pathologically diverse new species designated *C. pecorum* (Fukushi and Hirai, 1992; Kaltenboeck et al., 1993). This agent is not known to cause disease in humans.

Thus, *C. abortus* in cattle is predominantly associated with genital tract disease and mastitis. Exposure of pregnant women to *C. abortus* infected ruminants may lead to abortion or stillbirth (Hadley et al., 1992; Pospischil et al., 2002; Longbottom and Coulter, 2003; Baud et al., 2008).

Over the years, *Chlamydia* diagnosis in animals has improved (Sachse and Longbottom, 2013), and recent use of highly specific and sensitive nucleic amplification methods has also identified *C. psittaci*, albeit less frequently (Borel et al., 2006; Twomey et al., 2006; Kauffold et al., 2007; Pantchev et al., 2009; Kemmerling et al., 2009), and rarely *C. suis* in cattle (Teankum et al., 2007; Pantchev et al., 2009). Birds and swine are respectively, the main hosts for these two species. *C. psittaci* causes reproductive failure and respiratory disease in cattle, while the clinical significance of the occurrence of *C. suis* in cattle remains unclear (Longbottom and Coulter, 2003; Reinhold et al., 2011).

Chlamydiaceae infections in sheep and goats are caused by *C. abortus* and *C. pecorum*. Despite the existence of commercially available vaccines, *C. abortus*-induced lamb loss and enzootic abortion during the last third of gestation still remain the most common causes of reproductive failure in sheep-breeding countries all over the world. *C. abortus* causes major economic losses in affected flocks. Enzootic abortion in ewes (OEA) is a notifiable disease in Belgium and is also notifiable to the OIE. *C. pecorum* occasionally causes abortion in small ruminants. Especially in lambs, *C. pecorum* can induce, according to the subtype, pneumonia, polyarthritis, conjunctivitis, enteritis or clinically inapparent infections (Rodolakis et al., 1998).

The authors could not find any data on zoonotic transfer of *C. abortus* in Belgium; perhaps, because of unawareness by physicians, which is due to the absence of an efficient risk assessment.

It is difficult to conduct a zoonotic risk assessment, as epidemiological data on *C. abortus* in Belgium are lacking. Therefore, in the present study, the seroprevalence of *C. abortus* on Belgian sheep, goats and cattle farms is examined.

MATERIALS AND METHODS

Transversal seroepidemiological study in small and large ruminants

Sheep (n=958), goats (n=48) and cattle (1849) sera were provided by the biobanks of the following governmental institutions ARSIA (Association Régionale de Santé et d'Identification Animales, Ciney), DGZ (Dierengezondheidszorg Vlaanderen, Drongen) and CODA-CERVA (Veterinary and Agrochemical Research Centre, Brussels). For sheep and cattle, the examined herds were proportional to the number of herds in each Belgian province. The survey included 95 sheep herds from the list of volunteers enrolled in the Visna-Maedi certification program. For cattle herds (n=129), samples originated from the whole Belgian population. In addition, available sera of nine goat herds were serologically examined. The available sera of cattle were limited to animals ageing at least 24 months, while sheep and goat sera were not restricted by age category. None of the examined animals were vaccinated against *C. abortus*.

Sample collection and preparation were as follows: blood was collected by vene puncture during the winter of 2009-2010, incubated overnight at room temperature, centrifuged (325 x g, 4°C, 10 min), and serum was collected and stored at -20 °C. All sera were tested for the presence of antibodies against *C. abortus* using the same batch of the ID Screen™ *Chlamydia abortus* indirect multi-species ELISA (IDVET Innovative Diagnostics, Montpellier, France). The assay was performed according to the instructions of the manufacturer. Samples were tested at a dilution of 1/100. For each sample, the S/P was calculated, which is 100 times the OD₄₅₀ of the sample/mean value of the positive control OD₄₅₀. Samples presenting a S/P of greater than or equal to 60% were considered positive.

Statistics

All data were recorded in a MS Access® relational database. In sheep, the following variables were recorded at animal level: identity, disease status, probability of being sampled among the population herd of origin (Dohoo et al., 2010). At herd level size (two categories: ≤ and > 20 animals), disease status, geographical location, probability of being sampled and

province were recorded. In cattle, the same variables were recorded but four categories in herd size were considered (1-10, 11-50, 51-120 and >120 animals). At animal level, the age category was added and at herd level the production type (pure beef production versus other) was recorded.

Both at animal and herd levels, logistic regressions were used to assess apparent seroprevalences of *C. abortus* and to analyse potential multivariate relationships between risk factors (recorded variables) and individual disease status using STATA SE (StataCorp 4905 Lakeway Drive, College Station, Texas 77845, USA) also providing 95% confidence intervals (CI). Herd clustering was taken into account as random effect (Dohoo et al., 2010).

RESULTS

Transversal seroepidemiological study in sheep

From nine different Belgian regions, 958 sheep sera were tested (Table 1). The highest prevalence rate was found in Limburg (4.05%). The animals sampled in Antwerp, Brabant and Liège tested negative. The prevalence in the remaining five regions was rather low, ranging from 0.24% to 2.74%. Prevalence rates for all seropositive regions were statistically the same. The multivariate logistic regression model revealed a seroprevalence rate of 0.68% (95% confidence inter-

val (95 CI), 0.30-1.49) for the tested Belgian sheep population.

Fifteen of 95 (15.7%) sheep herds tested seropositive. The highest percentage of positive herds was found in Hainaut (Table 2). The *C. abortus* herd level prevalence was 6.15% (95% CI, 1.93-17.94), as calculated by the multivariate logistic regression model. Only herds with less than 50 sheep were seropositive. The *C. abortus* prevalence rate was higher in smaller herds (Table 3). However, taken all together, the *Chlamydia abortus* morbidity in the sheep was low.

Transversal seroepidemiological study in goats

Nine goat herds were tested, of which two herds with sample size ≥ 10 and seven herds with sample size <10 . None of the seven low-sample size herds tested seropositive. One of the two higher sample size herds tested seropositive. This herd, which was located in Luxembourg, revealed a high intraherd prevalence (52.9%; 9/17).

Transversal seroepidemiological study in cattle

From ten different Belgian regions, 1849 sera were tested (Table 4). The highest prevalence rate was found in Walloon Brabant (4.23%). The animals sampled in Limburg and Namur tested negative. The prevalence rate for the remaining seven regions ranged between 0.39% and 4.02%, without being statistically different

Table 1. *C. abortus* seroprevalence in sheep (N = 958).

Region	Sheep			
	Animals tested (N)	Prevalence (%)	95% CI Inferior limit (%)	95% CI Superior limit (%)
Antwerp	60	0.00	0.00	0.00
Brabant	134	0.00	NC	NC
Hainaut	100	0.79	0.36	6.56
Liège	70	0.00	NC	NC
Limburg	50	4.05	0.92	16.13
Luxembourg	82	2.74	0.37	17.53
Namur	61	0.24	0.01	3.83
East-Flanders	210	0.49	0.15	1.60
West-Flanders	191	0.93	0.22	3.85

Table 2. *C. abortus* seroprevalence in sheep herds (N = 95).

Region	Herds tested (N)	Prevalence (%)	Inferior limit (%)*	Superior limit (%)*
Antwerpen	8	0.0	0	-
Brabant	11	0.0	0	-
Hainaut	9	33.3	11.1	66.7
Liège	6	0.0	0	-
Limburg	9	22.2	5.6	57.9
Luxembourg	5	20.0	2.7	69.1
Namur	6	16.7	2.3	63.1
East-Flanders	25	16.0	6.1	35.7
West-Flanders	16	18.8	6.2	44.7

*Multivariate logistic regression with 95% confidence interval - Impossible to calculate the upper limit of the 95% confidence interval.

Table 3. Number of positive and negative sheep herds in function of herd size and minimum and maximum prevalences in positive herds.

Herd size	Herd status		Intraherd seroprevalence (%) for positive herds*		
	Negative	Positive	Minimum	Mean	Maximum
1 to 10	42	8	5.88	12.23	20.00
11 to 50	32	6	5.88	9.97	16.67
51 to 100	2	0	0.00	0.00	0.00
> 100	5	0	0.00	0.00	0.00

Table 4. *C. abortus* seroprevalence in cattle (N = 1849).

Region	Animals tested (N)	Prevalence (%)	Inferior limit (%)*	Superior limit (%)*
Antwerp	183	3.66	1.83	7.21
Hainaut	247	1.45	0.47	4.39
Liège	246	4.02	0.92	15.90
Limburg	83	0.00	0.00	8.74
Luxemburg	161	3.86	1.22	11.53
Namur	106	0.00	NC	NC
East-Flanders	356	0.41	0.10	1.67
Flemish Brabant	87	0.61	0.07	4.96
Walloon Brabant	30	4.23	2.67	6.63
West-Flanders	350	0.39	0.07	2.16

Table 5. *C. abortus* seroprevalence in cattle herds (N = 129).

Region	Herds tested (N)	Prevalence (%)	Inferior limit (%)*	Superior limit (%)*
Antwerp	14	35.71	15.70	62.37
Hainaut	16	25.00	9.71	50.82
Liège	16	12.50	3.14	38.60
Limburg	7	0.00	0.00	64.24
Luxemburg	9	22.22	5.60	57.90
Namur	8	0.00	0.00	68.62
East-Flanders	27	7.41	1.86	25.25
	7	14.29	1.97	58.06
Walloon Brabant	3	33.33	4.34	84.65
West-Flanders	22	9.09	2.28	29.96

Table 6. Number of positive and negative cattle herds in relation to herd size and minimum and maximum prevalences in positive herds.

Herd size	Herd status		Intraherd seroprevalence (%) for positive herds*		
	Negative	Positive	Minimum	Mean	Maximum
1 to 10	21	0	0.0	0.0	0.0
11 to 50	20	4	5.0	7.4	12.5
51 to 120	41	7	5.0	8.3	17.6
> 120	28	8	5.0	10.6	30.0

(Table 4). The prevalence rate was higher (albeit not statistically) in 24- to 48-months-old animals (prevalence 2.32%, 95% CI, 1.07-4.95), than in animals older than 48 months (prevalence 1.07%, 95% CI, 0.56-2.04). The prevalence rate was also higher (albeit not statistically) in beef cattle (prevalence 1.82%, 95% CI, 0.70-4.62), as compared to dairy or cross bred cattle (prevalence 1.59%, 95% CI, 0.72-3.46). For the cattle population, the multivariate logistic regression

model revealed a seroprevalence rate of 1.69% (95% CI, 0.91-3.10).

Nineteen of 129 (14.72%) cattle herds tested seropositive. Looking at herd level, the highest percentage of positive cattle herds was found in Antwerp (Table 5). The *C. abortus* prevalence on Belgian herd level was 14.7% (95% CI, 9.60-21.90), as calculated by the multivariate logistic regression model. Again, no significant differences were observed regarding

age or cattle breed. Interestingly, all positive herds consisted out of at least eleven animals (Table 6). Larger herds (11 to 50, 50 to 120 or <120 animals) were more frequently positive (albeit not significantly) and the number of positive animals per herd augmented with an increasing herd size (although not significantly) (Table 6). Morbidity due to *Chlamydia abortus* in cattle was rather low, but albeit 2.5 times higher than in sheep.

DISCUSSION

C. abortus is frequently isolated from ruminants and is responsible for abortion, infertility, keratoconjunctivitis, pneumonia, enteritis, mastitis and arthritis (Reinhold et al., 2011). Transmission occurs orally or sexually. *C. abortus* is a zoonotic pathogen. Several cases of abortion have been reported in pregnant women after contact with sheep (McKinlay et al., 1985; Herring et al., 1987; Longbottom and Coulter, 2003; Walder et al., 2005; Janssen et al., 2006) and goats (Pospischil et al., 2002; Meijer et al., 2004). Human infection is characterized by acute flu-like symptoms followed by abortion and in some cases renal failure, hepatic dysfunction and extensive intravascular coagulation resulting in death. No information is available on the *C. abortus* infection status in Belgian ruminants or on the risk for human health. The authors therefore conducted a seroprevalence study in cattle, sheep and goats. The ID Screen™ *Chlamydia abortus* indirect multi-species ELISA was used (ID-VET Innovative Diagnostics, Montpellier, France), since it is the only commercially available *C. abortus* ELISA, which uses microwells coated with a *C. abortus*-specific synthetic peptide from the major outer membrane protein (MOMP). The specificity of the test is 99.5%, while the sensitivity for small ruminants is expected to be 80% (difficult to find a large infected population) (P. Pourquier, personal communication, 2012). However, in cattle, the sensitivity is even more difficult to determine, because of very low confirmed *C. abortus* cases in cattle (P. Pourquier, personal communication, 2012).

Current results revealed a seroprevalence rate of 1.69% for the Belgian cattle population. Seroprevalence in positive herds was relatively low, with generally only 1 or 2 seropositive animals on 10 to 20 tested animals per herd. This may explain a much higher rate of seronegativity at herd level in herds for which < 10 animals were tested. Wilson et al., (2012) performed a study in Irish cattle (100 herds, 20 samples/herd) using a soluble chlamydial antigen (detergent treated *Chlamydia* elementary bodies) ELISA detecting antibodies against both *C. abortus* and *C. pecorum* and reported a seroprevalence rate of 6.04% at animal level. Their results are comparable to the low seroprevalence found in the present study. However, the prevalence at herd level is much higher in Irish herds (60%). The seropositive rate in Belgian cattle

herds in the current study could be an underestimation of the real level of infection of cattle in Belgium as the sensitivity of the ID VET ELISA could not yet be tested due to very few confirmed *C. abortus* cases in cattle. This might also explain the apparent absence of infected herds in the provinces of Limburg and Namur. Consequently, the human exposure rate may be underestimated. Seroprevalence studies on cattle in most other countries have reported higher infection rates at animal level (Reinhold et al., 2011). For instance, Wehrend et al. (2005) used a genus-specific enzyme-linked immunosorbent assay (ELISA). They found a seroprevalence of 41.5% in 445 dairy cows from 34 German farms, which is probably not abnormal as the ELISA detected antibodies against all members of the genus *Chlamydia*. Cavarani et al. (2001) examined 671 dairy cows of the Po Valley of Northern Italy using a commercial indirect ELISA (CHEKIT, Bommeli AG- IDEXX), and found a prevalence rate of 24.0%. These two studies focused only on dairy farms with fertility problems, which could explain the higher infection rates than in Belgian cattle, as this subpopulation has probably higher infection levels than the general population. Then again, results of Godin et al. (2008) suggest that *C. abortus* is rare in Swedish dairy cows with reproductive disorders. These authors examined 525 cows in 70 Swedish dairy herds by the use of both the CHEKIT ELISA and the commercial Pourquier *C. abortus* ELISA (Institut Pourquier, Montpellier, France). The Pourquier ELISA, in which a recombinant fragment of an 80–90 kDa polymorphic outer membrane protein as antigen is used, revealed a seroprevalence rate of 0.4% at animal level. The CHECKIT ELISA however, showed a seroprevalence of 28% (Godin et al., 2008). These and other experimental results indicate a lower specificity of the CHECKIT ELISA than of the Pourquier ELISA (Vretou et al., 2007; Godin et al., 2008; Wilson et al., 2009). The authors of the present study decided to use the Pourquier *C. abortus* ELISA. However, it has recently been removed from the market.

In the current study, serological examination of the sheep revealed a seroprevalence rate of 0.68%. As for cattle, the occurrence of *C. abortus* antibodies appears low in Belgian sheep as compared to most other reports on seroprevalence rates in non-vaccinated sheep: Switzerland (9.2 to 19.0%) (Borel et al., 2004, 2006; Blumer et al., 2012), Ireland (11%) (Markey et al., 1993) and Germany (15.1 to 94.0%) (Lenzko et al., 2011; Runge et al., 2012). Nevertheless, large variations in reported seroprevalence rates may in part be due to differences in sensitivity and/or specificity of serological tests used and/or to the size of the population under study. One of two Belgian goat herds that were examined at sample size ≥ 10 , tested seropositive, revealing an intraherd seroprevalence of 52.9% (9/17). The seroepidemiological study in goats should be expanded to a larger population.

It is noteworthy that in the UK, the number of reported cases in sheep is considerably higher in farms

holding more than 150 animals (47.6%) than in smaller herds (< 150 animals) (9.4%) (Longbottom et al., 2012). The authors of the present study obtained opposite results: no infection in flocks of larger size. Moreover, *C. abortus* morbidity in Belgian sheep seems to be low given the low prevalence in infected herds. This poses questions about the size of the examined population, which might have been too small. It is therefore possible that the rate of infection in sheep is underestimated. A study of management conditions and risk factors, as initially planned, could answer these questions. It is therefore regrettable, albeit understandable, that the sector was not very willing to provide an active contribution to this study, as it was difficult to collect enough sheep serum samples.

The obtained data suggest a limited infectivity for *C. abortus* in sheep and cattle, as the intra-herd seroprevalence in the positive herds was rather low. Wilson et al., (2012) observed the same. In the light of the current results, further research on the development of sensitive and specific *C. abortus* serological assays would be useful. These tests could also be helpful to define the actual exposure level for the human population at risk. In the future, it would also be worthwhile to perform molecular diagnoses on samples obtained from herds with clearly documented clinical data to get insight into the number of conjunctivitis, mastitis and reproductive failure cases in Belgium, where *C. abortus*, perhaps in conjunction with other pathogens, might be involved.

ACKNOWLEDGEMENTS

This study was funded by the Federal Public Service of Health, Food Chain Safety and Environment (convention RT 08/5 EMBACZON). Lizi Yin has a PhD fellowship from the China Scholarship Council (CSC grant; 01SC2812) and from the Special Research Fund of Ghent University (co-funding of the CSC grant).

REFERENCES

- Baud D., Regan L., Greub G. (2008). Emerging role of *Chlamydia* and *Chlamydia*-like organisms in adverse pregnancy outcomes. *Current Opinion in Infectious Diseases* 21, 70-76.
- Beer R., Bradford W., Hart R. (1982). Pregnancy complicated by psittacosis acquired from sheep. *British Medical Journal* 284, 1156-1157.
- Borel N., Doherr M., Vretou E., Psarrou E., Thoma R., Pospischil A. (2004). Seroprevalences for ovine enzootic abortion in Switzerland. *Preventive Veterinary Medicine* 65, 205-216.
- Blumer S., Moestl K., Krametter-Froetscher R., Hässig M., Pospischil A., Borel N. (2012). The seroconversion of *Chlamydia abortus* in sheep from the region of Vorarlberg before and after Alpine pasturing. *Schweizer Archiv für Tierheilkunde* 154, 13-17.
- Borel N., Thoma R., Spaeni P., Weilenmann R., Teankum K., Brugnera E., Zimmermann D.R., Vaughan L., Pospischil A. (2006). *Chlamydia*-related abortions in cattle from Graubünden, Switzerland. *Veterinary Pathology Online* 43, 702-708.
- Cavirani S., Cabassi C., Donofrio G., De Iaco B., Taddei S., Flammini C. (2001). Association between *Chlamydia psittaci* seropositivity and abortion in Italian dairy cows. *Preventive Veterinary Medicine* 50, 145-151.
- Dohoo I. Martin W., Stryhn H. (2010). Mixed models for discrete data. In: S.M. McPike (ed.). *Veterinary Epidemiologic Research*. Second edition. VER Inc., Prince Edwards Island, Canada, 579-606.
- Everett K.D., Bush R.M., Andersen A.A. (1999). Emended description of the order *Chlamydiales*, proposal of *Parachlamydiaceae* fam. nov. and *Simkaniaceae* fam. nov., each containing one monotypic genus, revised taxonomy of the family *Chlamydiaceae*, including a new genus and five new species, and standards for the identification of organisms. *International Journal of Systematic Bacteriology* 49, 415-440.
- Fukushi H., Hirai K. (1992). Proposal of *Chlamydia pecorum* sp. nov. for *Chlamydia* strains derived from ruminants. *International Journal of Systematic Bacteriology* 42, 306-308.
- Godin A.C., Björkman C., Englund S., Johansson K.E., Niskanen R., Alenius S. (2008). Investigation of *Chlamydophila* spp. in dairy cows with reproductive disorders. *Acta Veterinaria Scandinavica* 50, 39.
- Hadley K., Carrington D., Frew C., Gibson A., Hislop W. (1992). Ovine chlamydiosis in an abattoir worker. *Journal of Infection* 25, 105-109.
- Herring A., Anderson I., McClenaghan M., Inglis N., Williams H., Matheson B.A., West C.P., Rodger M., Brett P.P. (1987). Restriction endonuclease analysis of DNA from two isolates of *Chlamydia psittaci* obtained from human abortions. *British Medical Journal (Clinical Research ed.)* 295, 1239.
- Janssen M.J., Van de Wetering K., Arabin B. (2006). Sepsis due to gestational psittacosis: A multidisciplinary approach within a perinatal center--review of reported cases. *International Journal of Fertility and Women's Medicine* 51, 17-20.
- Kaltenboeck B., Kousoulas K., Storz J. (1993). Structures of and allelic diversity and relationships among the major outer membrane protein (*ompA*) genes of the four *Chlamydia* species. *Journal of Bacteriology* 175, 487-502.
- Kaltenboeck B., Hehnen H.R., Vaglenov A. (2005). Bovine *Chlamydophila* spp. infection: Do we underestimate the impact on fertility? *Veterinary Research Communications* 29, 1-15.
- Kauffold J., Henning K., Bachmann R., Hotzel H., Melzer F. (2007). The prevalence of *Chlamydiae* of bulls from six bull studs in Germany. *Animal Reproduction Science* 102, 111-121.
- Kemmerling K., Müller U., Mielenz M., Sauerwein H. (2009). *Chlamydophila* species in dairy farms: Polymerase chain reaction prevalence, disease association, and risk factors identified in a cross-sectional study in western Germany. *Journal of Dairy Science* 92, 4347-4354.
- Kuo C., Stephens R. (2011). Family I. *Chlamydiaceae*. In: Whitman, B. William (editors). *Bergey's Manual of Systematic Bacteriology*. Second edition, Springer Science/Business Media, New York, USA, p. 845.

- Lenzko H., Moog U., Henning K., Lederbach R., Diller R., Menge C., Sachse K., Sprague L.D. (2011). *BMC Veterinary Research* 16, 7:29.
- Longbottom D. and L.J. Coulter. (2003). Animal Chlamydiosis and zoonotic implications. *Journal of Comparative Pathology* 128, 217-244.
- Longbottom D., Entrican G., Wheelhouse N., Brough H., Milne C. (2012). Evaluation of the impact and control of enzootic abortion of ewes. *The Veterinary Journal* 195, 257-259.
- Markey B., McNulty M., Todd D. (1993). Comparison of serological tests for the diagnosis of *Chlamydia psittaci* infection of sheep. *Veterinary Microbiology* 36, 233-252.
- McKinlay A., White N., Buxton D., Inglis J., Johnson F., Kurtz J., Brettle R.P. (1985). Severe *Chlamydia psittaci* sepsis in pregnancy. *Quarterly Journal of Medicine* 57, 689-696.
- McNutt S.H., Waller E.F. (1940). Sporadic bovine encephalomyelitis. *The Cornell Veterinarian* 30, 437-448.
- Meijer A., Brandenburg A., Vries J., Beentjes J., Roholl P., Dercksen D. (2004). *Chlamydomydia abortus* infection in a pregnant woman associated with indirect contact with infected goats. *European Journal of Clinical Microbiology and Infectious Diseases* 23, 487-490.
- Pantchev A., Sting R., Bauerfeind R., Tyczka J., Sachse K. (2009). New real-time PCR tests for species-specific detection of *Chlamydomydia psittaci* and *Chlamydomydia abortus* from tissue samples. *The Veterinary Journal* 181(2), 145-150.
- Pospischil A., Thoma R., Hilbe M., Grest P., Gebbers J. (2002). Abortion in woman caused by caprine *Chlamydomydia abortus* (*Chlamydia psittaci* serovar 1). *Swiss Medical Weekly* 132, 64-66.
- Reinhold P., Sachse K., Kaltenboeck B. (2011). *Chlamydiaceae* in cattle: Commensals, trigger organisms, or pathogens? *The Veterinary Journal* 189, 257-267.
- Rodolakis A., Salinas J., Papp J. (1998). Recent advances on ovine chlamydial abortion. *Veterinary Research* 29, 275-288.
- Runge M., Binder A., Schotte U., Ganter M. (2012). Investigations concerning the prevalence of *Coxiella burnetii* and *Chlamydia abortus* in sheep in correlation with management systems and abortion rate in Lower Saxony in 2004. *Berliner und Münchener Tierärztliche Wochenschrift* 6, 138-143.
- Sachse K., Hotzel H., Slickers P., Ellinger T., Ehrlich R. (2005). DNA microarray-based detection and identification of *Chlamydia* and *Chlamydomydia spp.* *Molecular and Cellular Probes* 19, 41-50.
- Sachse K., Longbottom D. (2013). Enzootic abortion in ewes (ovine chlamydiosis). In: OIE Biological Standards Commission (eds.). *The Manual of Diagnostic Tests and Vaccines for Terrestrial Animals*. 7th Edition, Volume 2, p 1-9.
- Teankum K., Pospischil A., Janett F., Brugnera E., Hoelzle L., Hoelzle K., Weilenmann R., Zimmermann D.R., Gerber A., Polkinghorne A., Borel N. (2007). Prevalence of *Chlamydiae* in semen and genital tracts of bulls, rams and bucks. *Theriogenology* 67, 303-310.
- Twomey K.M., Rao S.V., Becker R.C. (2006). Proinflammatory, immunomodulating, and prothrombotic properties of anemia and red blood cell transfusions. *Journal of Thrombosis and Thrombolysis* 21, 167-174.
- Vretou E., Radouani F., Psarrou E., Kritikos I., Xylouri E., Mangana O. (2007). Evaluation of two commercial assays for the detection of *Chlamydomydia abortus* antibodies. *Veterinary Microbiology* 123, 153-161.
- Walder G., Hotzel H., Brezinka C., Gritsch W., Tauber R., Würzner R., Ploner F. (2005). An unusual cause of sepsis during pregnancy: recognizing infection with *Chlamydomydia abortus*. *Obstetrics & Gynecology* 106, 1215-1217.
- Wehrend A., Failing K., Hauser B., Jäger C., Bostedt H. (2005). Production, reproductive, and metabolic factors associated with chlamydial seropositivity and reproductive tract antigens in dairy herds with fertility disorders. *Theriogenology* 63, 923-930.
- Wilson K., Livingstone M., Longbottom D. (2009). Comparative evaluation of eight serological assays for diagnosing *Chlamydomydia abortus* infection in sheep. *Veterinary Microbiology* 135, 38-45.
- Wilson K., Sammin D., Harmeyer S., Nath M., Livingstone M., Longbottom D. (2012). Seroprevalence of chlamydial infection in cattle in Ireland. *The Veterinary Journal* 193, 583-585.